

# Impact of in vitro fertilization-embryo transfer on mother-toinfant transmission in women with chronic HBV infection

```
Wei Yi<sup>1</sup> Minghui Li<sup>2,3</sup> Fangfang Sun<sup>2</sup> Huihui Lu<sup>2</sup> Zhan Zeng<sup>3</sup> Xiaoyue Bi<sup>2</sup>
Liu Yang<sup>2</sup> | Yanjie Lin<sup>3</sup> | Xiuzhen Cao<sup>1</sup> | Yuhong Hu<sup>1</sup> | Mingfang Zhou<sup>1</sup> | Lu Zhang<sup>2</sup> |
Yao Lu<sup>2</sup> | Gang Wan<sup>4</sup> | Yao Xie<sup>2,3</sup> ©
```

Correspondence

and

**Funding information** 

**Abstract** Background and Aims:

Methods:

Results:

Conclusion:

**Handling Editor:** 

#### 1 | INTRODUCTION

1

6 6–8

10

#### 2.3 | Prophylactic immunization of newborn

μ

= = ± 10

X

#### 2.4 | Outcome measures

## 2.5 | Statistical analysis

 $\pm$ 

val

#### 3 | RESULTS

# 3.1 | Enrollment and clinical characteristics of pregnant women

Values	CHB group ( = 224)	Control group ( = 74)	High viral load -value group ( = 95)	Low viral load group $(=129)$		Treated group -value ( = 72)	Untreated group $(=23)$	-value
+1	+1	+1	+1	+1	.102	+1	+1	
+1	+1							

14-16

WILEY-	Liver	
• • • • • •	INTERNATIONAL	

ILLI	IN	TERN	ATIO	NAL			SHE	K		
-value			.240	.020	.633	1.000	.160	I	I	
Untreated group ( = 26)	+1	+I	+1		+1					
Treated group ( = 87)	+1	+I	+1		+1					
-value		.726		998.	.314	.246	.028		1	
Low viral load group ( = 169)	+1	+I	+I		+1					
High viral load group $(=113)$	+1	+1	+1		+1					
-value	.167	.083	.212		.064	1.000	ı	ı	ı	
Control group $(=93)$	+1	+I	+1		+1		I	I	1	
CHB group ( = 282)	+1	+I	+I		+1					
	+1		+1		+1					
Characteristic		+I					+	+		

TABLE 3

## 4 | DISCUSSION

.21 22-24 µ

.12

12

•

CONFLICT OF INTEREST

**ETHICS APPROVAL** 

CLINICAL TRIAL NUMBER

ORCID

D D

REFERENCES

**FUNDING INFORMATION** 

. 2017;77:e12636.

. 2011;26:1860e7.

varicocele.